

2

OFFICE OF NAVAL RESEARCH

Contract N00014-85-K-0772 Work Unit No. 4326-808

TECHNICAL REPORT NO. 139

A Ferracarborane Analogue to [Fp]⁻. Synthesis and Reactions of [closo-3,3-(CO)₂-3,1,2-FeC₂B₉H₁₁]²-

by
Sharon S. Lee, Carolyn Knobler, and M. Frederick Hawthorne*



Prepared for Publication

in

Organometallics

University of California at Los Angeles
Department of Chemistry and Biochemistry
Los Angeles, California 90024

April, 1991

A-1

Reproduction in whole or in part is permitted for any purpose of the United States Government

This document has been approved for public release and sale; its distribution is unlimited

91-07105

REPORT DOCUMENTATION PAGE									
1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED				1b. RESTRICTIVE MARKINGS					
2a. SECURITY CLASSIFICATION AUTHORITY				3. DISTRIBUTION / AVAILABILITY OF REPORT					
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE				Distribution list enclosed					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) 139				5. MONITORING ORGANIZATION REPORT NUMBER(S)					
UCLA		Biochem.	6b. OFFICE SYMBOL (If applicable)	7a NAME OF MONITORING ORGANIZATION Office of Naval Research					
405 Hil Los Ang	oung Hall Lgard Aven geles, CA	iue 90024	7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Arlington, VA 22217						
Ba. NAME OF I ORGANIZA' Office	TION	nsoring Research	8b OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER					
Bc. ADDRESS (C	ity, State, and	ZIP Code)	<u> </u>	10 SOURCE OF	FUNDING NUMBER	S			
				PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO		WORK UNIT ACCESSION NO	
11. TITLE (Inclu	11. TITLE (Include Security Classification) A Ferracarborane Analogue to [Fp]. Synthesis and Reactions of [closo-3,3-(CO)2-3,1,2-FeC2B9H11]2								
12 PERSONAL	AUTHOR(S) Share	on S. Lee, Car	rolyn Knobler, a	nd M. Freder	rick Hawthor	ne			
13a TYPE OF REPORT 13b TIME COVERED FROM TO TO				14 DATE OF REPORT (Year, Month, Day) 15 PAGE COUNT 7-24-91					
16 SUPPLEMENTARY NOTATION To be published in Organometallics									
17	COSATI	CODES	18 SUBJECT TERMS (C	ontinue on revers	e if necessary and	identi	fy by bloci	k number)	
FIELD	GROUP	SUB-GROUP							
			carbonyl	borane, dicarbollide, iron, iron					
Reduction of [closo-3,3,3-(CO) ₃ -3,1,2-FeC ₂ B ₂ H ₁₁] (1) with 2 equiv of sodium naphthalide affords the high-yield synthesis of a formal iron(0) ferracarborane dianion [closo-3,3-(CO) ₂ -3,1,2-FeC ₂ B ₂ H ₁₁] ²⁻ (3), as monitored by ¹¹ B NMR spectroscopy. Complex 3 serves as a nucleophile in a variety of alkylation, acylation, and metalation reactions to yield ferracarborane anions of the type [closo-3,3-(CO) ₂ -3-L-3,1,2-FeC ₂ B ₂ H ₁₁] ¹⁻ (4, L ⇒ CH ₃ ; 5, L = CH ₂ C ₆ H ₆ ; 6, L = COCH ₃ ; 7, L = SnC ₆ H ₆). The molecular structure of [N(C ₂ H ₆) ₄][7] has been determined by X-ray diffraction techniques. Compound 7 crystallized in the triclinic space group P½ with a = 11.209 (1) Å, b = 13.026 (1) Å, c = 13.595 (1) Å, a = 79.959 (3)°, β = 84.143 (3)°, γ = 66.060 (2)°, V = 1785 ų, and Z = 2. In situ reactions of 3 with allyl and methylallyl chloride followed by subsequent loss of CO induced by ultraviolet radiation, resulted in the isolation of the corresponding η³-allyl derivatives, [closo-3-CO-3-(η³-CH ₂ CRCH ₂)-3,1,2-FeC ₂ B ₂ H ₁₁] (8, R = H; 9, R = CH ₃). The structure of 8 as the PPN* salt was also elucidated by single-crystal X-ray diffraction and found to crystallize in the monoclinic space group P2 ₁ /m with a = 9.1250 (7) Å, b = 25.011 (2) Å, c = 18.517 (2) Å, β = 91.328 (3)°, V = 4225 ų, and Z = 4 (two crystallographically different half-anions and one cation in the asymmetric unit). Migratory 20 DISTRIBUTION/AVAILABILITY OF ABSTRACT DTIC USERS									
	F RESPONSIBLE		RPT LIDTIC USERS	22b TELEPHONE (Include Area Code) 22c OFFICE SYMBOL					
M. F. Hawthorne (213)825-7378									

19. Abstract (cont.)

insertion of alkyls was demonstrated in the preparation of $\{closo\text{-}3\text{-}CO\text{-}3\text{-}COCH_5\text{-}3\text{-}P(CH_2)_3\text{-}3,1,2\text{-}FeC_2B_2H_{11}\}$ (10) by heating 4 and excess $P(CH_3)_3$ in THF for 7 days at the reflux temperature. A single-crystal X-ray diffraction study of [PPN][10] is reported. Complex 10 crystallized in the triclinic space group P^{I} with a=10.086 (4) Å, b=15.390 (6) Å, c=17.027 (7) Å, $\alpha=112.636$ (8)°, $\beta=96.647$ (8)°, $\gamma=100.206$ (9)°, V=2351 ų, and Z=2. The closo 12-vertex icosahedral geometry composed of a polyhedral FeC_2B_3 - d_6 framework and pseudooctahedral coordination exhibited by the iron atom are common structural features displayed by all three of the ferracarboranes that were characterized crystallographically.

A Ferracarborane Analogue to [Fp]⁻. Synthesis and Reactions of [closo-3,3-(CO)₂-3,1,2-FeC₂B₉H₁₁]²⁻

Sharon S. Lee, Carolyn B. Knobler, and M. Frederick Hawthorne*

Department of Chemistry and Biochemistry, University of California, Los Angeles, Los Angeles, California 90024

Received July 6, 1990

Reduction of [closo-3,3,3-(CO)₃-3,1,2-FeC₂B₉H₁₁] (1) with 2 equiv of sodium naphthalide affords the high-yield synthesis of a formal iron(0) ferracarborane dianion [closo-3,3-(CO)₂-3,1,2-FeC₂B₉H₁₁]²⁻ (3), as monitored by ¹¹B NMR spectroscopy. Complex 3 serves as a nucleophile in a variety of alkylation, acylation, and metalation reactions to yield ferracarborane anions of the type [closo-3,3-(CO)₂-3-L-3,1,2-FeC₂B₉H₁₁]⁻ (4, L = CH₃; 5, L = CH₂C₆H₅; 6, L = COCH₃; 7, L = SnC₆H₅). The molecular structure of [N(C₂H₅)₄][7] has been determined by X-ray diffraction techniques. Compound 7 crystallized in the triclinic space group P\bar{1} with a = 11.209 (1) Å, b = 13.026 (1) Å, c = 13.595 (1) Å, \alpha = 79.959 (3)^{\alpha}, \beta = 84.143 (3)^{\alpha}, \gamma = 66.060 (2)^{\alpha}, V = 1785 Å³, and Z = 2. In situ reactions of 3 with allyl and methylallyl chloride followed by subsequent loss of CO induced by ultraviolet radiation, resulted in the isolation of the corresponding \(\eta^3\text{-allyl derivatives}\), [closo-3-CO-3-(\eta^3\text{-CH}_2CRCH_2)-3,1,2\text{-FeC}_2B₉H₁₁]⁻ (8, R = H; 9, R = CH₃). The structure of 8 as the PPN⁺ salt was also elucidated by single-crystal X-ray diffraction and found to crystallize in the monoclinic space group P2₁/m with a = 9.1250 (7) Å, b = 25.011 (2) Å, c = 18.517 (2) Å, \beta = 91.328 (3)^{\alpha}, V = 4225 Å³, and Z = 4 (two crystallographically different half-anions and one cation in the asymmetric unit). Migratory insertion of alkyls was demonstrated in the preparation of [closo-3-CO-3-COCH₃-3-P(CH₃)₃-3,1,2-FeC₂B₉H₁₁]⁻ (10) by heating 4 and excess P(CH₃)₃ in THF for 7 days at the reflux temperature. A single-crystal X-ray diffraction study of [PPN][10] is reported. Complex 10 crystallized in the triclinic space group P\bar{1} with a = 10.086 (4) Å, b = 15.390 (6) Å, c = 17.027 (7) Å, \alpha = 112.636 (8)^{\alpha}, \beta = 96.647 (8)^{\alpha}, \gamma = 100.206 (9)^{\alpha}, V = 2351 Å³, and

01/1113/86/2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies		No. Copies
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 2770	1 9
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 1911	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1